

## PREVALENCE OF DIABETES MELLITUS AMONG ADOLESCENTS, RISK FACTORS, AND IMPROVEMENT OF MEDICAL AND SOCIAL CARE

Orifova Munisa

Second-year Master's student, Tashkent State Medical University

Professor M.B. Mamatkulov

### **Abstract**

Diabetes mellitus among adolescents has become an increasingly important global public health concern due to its rising prevalence and long-term health consequences. Once considered predominantly an adult disease, diabetes is now frequently diagnosed during adolescence, driven by a combination of genetic predisposition, lifestyle changes, obesity, and socioeconomic factors. This scientific analytical study aims to examine the prevalence of diabetes mellitus among adolescents, identify major risk factors, and evaluate approaches for improving medical and social care. The analysis is based on a review of international epidemiological data, clinical studies, and public health reports. The findings indicate a steady increase in both type 1 and type 2 diabetes among adolescents, with type 2 diabetes showing particularly rapid growth in recent years. Key risk factors include obesity, physical inactivity, unhealthy dietary habits, family history of diabetes, and psychosocial stress. In addition, limited access to healthcare services and inadequate social support significantly affect disease management and outcomes. The study highlights the importance of integrated care models that combine medical treatment with psychological, family, school-based, and community-level interventions. Strengthening preventive strategies, early screening programs, and multidisciplinary care systems is essential for reducing disease burden and improving long-term health outcomes among adolescents with diabetes mellitus.

### **Key words**

Diabetes mellitus, adolescents, prevalence, risk factors, obesity, lifestyle factors, medical care, social care, prevention, public health.

### **Introduction**

Diabetes mellitus (DM) has emerged as one of the most significant global public health challenges of the 21st century, affecting individuals across all age groups. Traditionally considered a disease of adulthood, diabetes is now increasingly diagnosed among adolescents, signaling a worrying epidemiological shift. The rising prevalence of diabetes mellitus among adolescents is closely linked to rapid changes in lifestyle, nutrition patterns, urbanization, and socioeconomic conditions. This trend poses serious medical, psychological, and social consequences, as early-onset diabetes is associated with long-term complications, reduced quality of life, and increased healthcare burdens. Globally, the incidence and prevalence of both type 1 diabetes mellitus (T1DM) and type 2 diabetes mellitus (T2DM) among adolescents have shown a steady upward trajectory over the past decades. While T1DM remains the most common form in this age group due to autoimmune destruction of pancreatic  $\beta$ -cells, the prevalence of T2DM among adolescents has risen dramatically, particularly in low- and middle-income countries. This increase is largely attributed to childhood obesity, physical inactivity, unhealthy dietary habits, and genetic susceptibility. According to international health organizations, the growing burden of adolescent diabetes represents a critical challenge for healthcare systems, requiring early detection, effective management, and comprehensive preventive strategies.

Adolescence is a particularly vulnerable developmental period characterized by rapid physical growth, hormonal changes, and psychological transformation. These biological and psychosocial factors significantly influence metabolic regulation and insulin sensitivity, increasing susceptibility to glucose metabolism disorders. Moreover, adolescents often



demonstrate limited adherence to medical recommendations, irregular health-seeking behavior, and insufficient awareness of disease-related risks. As a result, diabetes diagnosed during adolescence frequently remains poorly controlled, leading to early microvascular and macrovascular complications such as retinopathy, nephropathy, neuropathy, and cardiovascular disorders. Risk factors for diabetes mellitus among adolescents are multifactorial and interrelated, encompassing genetic, environmental, behavioral, and social determinants. Family history of diabetes, obesity, sedentary lifestyle, excessive consumption of high-calorie and processed foods, and psychological stress are among the most prominent contributors. In addition, socioeconomic inequalities, limited access to healthcare services, inadequate health education, and cultural factors further exacerbate the risk and hinder timely diagnosis and effective disease management. Understanding these risk factors is essential for developing targeted prevention programs and personalized intervention strategies. Beyond its medical implications, diabetes mellitus among adolescents has profound social and psychological consequences. The diagnosis often affects academic performance, social integration, emotional well-being, and self-esteem. Adolescents with diabetes may experience stigma, anxiety, depression, and difficulties in peer relationships, which can negatively impact disease self-management and long-term outcomes. Furthermore, the chronic nature of diabetes places a significant burden on families and caregivers, emphasizing the need for integrated medical and social support systems.

Improvement of medical and social care for adolescents with diabetes is a key priority in addressing this growing health concern. Effective management requires a multidisciplinary approach involving pediatric endocrinologists, primary care physicians, nurses, psychologists, nutritionists, social workers, and educators. Early screening programs, school-based health interventions, patient-centered education, and community engagement play a crucial role in enhancing disease awareness, promoting healthy behaviors, and ensuring continuity of care. Advances in medical technologies, such as continuous glucose monitoring systems and insulin delivery devices, have also improved glycemic control; however, their accessibility remains limited in many regions. A comprehensive scientific analysis of the prevalence of diabetes mellitus among adolescents, associated risk factors, and existing medical and social care mechanisms is of paramount importance. Such an analysis can contribute to evidence-based policy development, optimization of healthcare services, and implementation of effective preventive measures. This article aims to examine current trends in adolescent diabetes prevalence, analyze key risk factors, and explore strategies for improving medical and social care to mitigate the long-term impact of diabetes mellitus on adolescent health and societal well-being.

### Literature review

The growing prevalence of diabetes mellitus among adolescents has attracted increasing attention from researchers and public health organizations worldwide. Numerous epidemiological, clinical, and social studies have examined trends in adolescent diabetes, associated risk factors, and the effectiveness of existing medical and social care systems. This section reviews key findings from international and regional literature, highlighting current knowledge gaps and research directions.

A substantial body of literature indicates a continuous rise in the prevalence of diabetes mellitus among adolescents over the past several decades. According to reports from the International Diabetes Federation (IDF) and the World Health Organization (WHO), type 1 diabetes mellitus (T1DM) remains the most common form among adolescents; however, type 2 diabetes mellitus (T2DM) has shown a particularly rapid increase, especially in urbanized and developing regions. Studies conducted in North America, Europe, and Asia demonstrate annual increases in T1DM incidence ranging from 2% to 5%, suggesting a strong influence of environmental and lifestyle-related factors in addition to genetic predisposition. Several population-based studies emphasize that the prevalence of adolescent diabetes varies



significantly across regions and socioeconomic contexts. High-income countries report better diagnostic coverage and disease registration, while low- and middle-income countries often underestimate the true prevalence due to limited screening programs and insufficient healthcare access. Researchers highlight that adolescent diabetes is no longer a rare condition and should be regarded as a major public health concern requiring coordinated global and national responses.

The literature consistently identifies adolescent diabetes mellitus as a multifactorial disease influenced by genetic, biological, behavioral, and social determinants. Genetic susceptibility, particularly a family history of diabetes, has been strongly associated with increased risk, especially for T2DM. However, recent studies emphasize that genetic predisposition alone cannot explain the rapid rise in prevalence, pointing instead to modifiable environmental factors. Obesity and overweight status are among the most frequently cited risk factors for adolescent T2DM. Numerous studies demonstrate a strong correlation between increased body mass index (BMI), insulin resistance, and impaired glucose metabolism in adolescents. Sedentary behavior, excessive screen time, and reduced physical activity have been identified as major contributors to obesity-related diabetes risk. In addition, dietary patterns characterized by high intake of sugar-sweetened beverages, fast food, and processed products significantly increase the likelihood of developing diabetes during adolescence. Hormonal and physiological changes during puberty also play a critical role in diabetes risk. Research indicates that puberty-related insulin resistance can exacerbate underlying metabolic abnormalities, particularly in genetically predisposed individuals. This transitional period is therefore considered a high-risk window for the onset of glucose metabolism disorders.

Beyond biological factors, a growing number of studies emphasize the importance of psychosocial and socioeconomic determinants in adolescent diabetes. Low socioeconomic status, limited parental education, and poor access to healthcare services have been consistently associated with delayed diagnosis, poor glycemic control, and increased complication rates. Adolescents from disadvantaged backgrounds often face barriers to healthy nutrition, physical activity, and continuous medical care. Psychological factors such as chronic stress, depression, anxiety, and low self-esteem have also been linked to increased diabetes risk and poor disease management. Adolescents with diabetes frequently experience emotional distress related to disease burden, fear of complications, and social stigma. Several qualitative studies report that psychosocial stress negatively affects treatment adherence, leading to suboptimal outcomes.

The medical management of adolescent diabetes has evolved significantly over recent decades. Advances in insulin therapy, glucose monitoring technologies, and evidence-based clinical guidelines have improved disease outcomes in many settings. Studies highlight the effectiveness of intensive insulin therapy and continuous glucose monitoring systems in achieving better glycemic control and reducing acute complications. However, the literature also points to persistent challenges in adolescent diabetes management. Adolescents often demonstrate lower adherence to treatment regimens compared to younger children and adults. Factors such as fear of injections, peer pressure, lack of disease awareness, and inadequate family support contribute to poor adherence. Researchers emphasize the need for age-specific, patient-centered approaches that consider the developmental and psychosocial characteristics of adolescents.

In recent years, increasing attention has been given to the role of social and community-based care in managing adolescent diabetes. Studies suggest that integrated care models combining medical treatment with psychological counseling, family education, and school-based interventions significantly improve health outcomes. School health programs, in particular, have been shown to enhance disease awareness, reduce stigma, and support daily diabetes management. Family involvement is widely recognized as a critical component of effective diabetes care. Literature indicates that adolescents with strong family support demonstrate better glycemic control, higher treatment adherence, and improved quality of life. Conversely, family



conflict and lack of parental engagement are associated with poor outcomes. Community-level interventions, including public health campaigns and peer support groups, have also shown promising results. These initiatives help promote healthy lifestyles, encourage early screening, and provide emotional support for adolescents living with diabetes.

Despite the extensive body of research on adolescent diabetes, several gaps remain. Many studies focus predominantly on clinical aspects, while social, cultural, and psychological dimensions are often underexplored, particularly in low-resource settings. Additionally, there is limited evidence on the long-term effectiveness of integrated medical and social care models in diverse populations. Researchers also note a lack of region-specific data, especially in Central Asia and other developing regions, where healthcare infrastructure and social support systems differ significantly from those in high-income countries. This gap underscores the need for localized studies to inform context-appropriate interventions and policy development. The reviewed literature demonstrates that diabetes mellitus among adolescents is a complex and growing global health issue driven by a combination of biological, behavioral, and social factors. While significant progress has been made in medical management, the effectiveness of care is closely linked to social support, psychological well-being, and healthcare accessibility. A comprehensive approach integrating medical treatment with social and preventive strategies is essential to address the rising burden of adolescent diabetes and improve long-term health outcomes.

### Research discussion

The findings discussed in this study highlight diabetes mellitus among adolescents as a rapidly escalating public health issue with complex medical, behavioral, and social dimensions. The increasing prevalence observed across different regions confirms trends reported in previous international studies and underscores the urgent need for comprehensive intervention strategies. The discussion integrates epidemiological patterns, identified risk factors, and implications for improving medical and social care systems. One of the most significant observations is the steady rise in diabetes prevalence among adolescents, particularly type 2 diabetes mellitus. This shift reflects broader global changes in lifestyle, nutrition, and physical activity patterns. Urbanization, increased consumption of energy-dense foods, and reduced physical activity have collectively contributed to higher rates of obesity, which remains a dominant modifiable risk factor. These findings are consistent with prior research emphasizing the strong association between adolescent obesity, insulin resistance, and early onset of metabolic disorders. The role of puberty-related physiological changes is also critical in understanding adolescent diabetes. Hormonal fluctuations during adolescence naturally reduce insulin sensitivity, which can exacerbate underlying metabolic vulnerabilities. When combined with obesity and genetic predisposition, these physiological changes significantly increase diabetes risk. This reinforces the importance of early screening and targeted preventive measures during the pubertal period, particularly for high-risk groups.

Genetic and family-related factors continue to play an important role in diabetes development. Adolescents with a family history of diabetes demonstrate higher susceptibility, supporting existing evidence that genetic predisposition interacts with environmental influences. However, the rapid increase in prevalence suggests that genetic factors alone cannot account for current trends. Instead, the interaction between heredity and lifestyle-related risk factors appears to be the primary driver of disease onset in this population. Beyond biological determinants, the discussion highlights the substantial influence of psychosocial and socioeconomic factors on both disease development and management. Adolescents from lower socioeconomic backgrounds often experience limited access to healthcare services, inadequate health education, and barriers to healthy lifestyle choices. These conditions not only increase the risk of diabetes but also contribute to delayed diagnosis and suboptimal disease control. The findings emphasize



that diabetes among adolescents should not be viewed solely as a medical condition but as a socially influenced disease requiring broader systemic interventions.

Psychological challenges associated with adolescent diabetes emerged as a critical concern. Adolescents frequently experience emotional stress, anxiety, and reduced self-esteem following diagnosis, which can negatively affect treatment adherence and glycemic control. Peer pressure and fear of social stigma further complicate disease management, particularly in school settings. These findings align with previous studies demonstrating that psychological well-being is closely linked to effective self-management and long-term outcomes in chronic diseases. Medical management strategies for adolescent diabetes have improved significantly due to advances in insulin therapy, glucose monitoring technologies, and standardized clinical guidelines. However, the discussion reveals persistent challenges related to treatment adherence and continuity of care. Adolescents often struggle to follow complex treatment regimens, especially when adequate education and psychosocial support are lacking. This highlights the need for age-appropriate, patient-centered care models that account for the developmental and emotional characteristics of adolescents. The importance of integrated medical and social care is strongly supported by the findings. Multidisciplinary care approaches involving healthcare professionals, psychologists, educators, and social workers have been shown to improve glycemic control, treatment adherence, and quality of life. School-based health programs play a particularly valuable role by providing daily support, increasing disease awareness, and reducing stigma among peers. These programs also create opportunities for early identification of symptoms and timely referral to medical services.

Family involvement remains a cornerstone of effective diabetes management in adolescents. Supportive family environments are associated with better disease outcomes, while family conflict or lack of engagement often leads to poor adherence and increased complication risk. The discussion emphasizes the need for family-centered education programs that empower parents and caregivers with the knowledge and skills required to support adolescents in managing their condition effectively. From a public health perspective, the findings suggest that preventive strategies must extend beyond clinical settings. Community-based interventions promoting healthy nutrition, physical activity, and regular health screening are essential for reducing the incidence of adolescent diabetes. Public awareness campaigns and peer support initiatives can help address misconceptions, reduce stigma, and encourage early health-seeking behavior. Such interventions are particularly important in regions with limited healthcare infrastructure. Despite its contributions, this study has certain limitations that should be acknowledged. The discussion relies on existing literature and secondary data, which may not fully capture region-specific variations in disease prevalence and care accessibility. Additionally, differences in diagnostic criteria, data collection methods, and healthcare systems across studies may affect comparability. These limitations highlight the need for longitudinal and regionally focused research to better understand local determinants of adolescent diabetes and evaluate the effectiveness of integrated care models. The discussion underscores that diabetes mellitus among adolescents is a multifaceted health challenge driven by the interaction of biological, behavioral, and social factors. While medical advancements have improved disease management, sustainable progress depends on strengthening social support systems, enhancing health education, and addressing socioeconomic inequalities. A holistic approach that integrates medical treatment with psychological, family, school, and community-based interventions is essential to reduce disease burden, prevent complications, and improve long-term outcomes for adolescents living with diabetes.

## Conclusion

Diabetes mellitus among adolescents represents a growing global health concern with significant medical, psychological, and social implications. The increasing prevalence of both



type 1 and type 2 diabetes highlights the impact of lifestyle changes, obesity, and socioeconomic factors on adolescent health. Evidence indicates that adolescence is a critical period during which biological vulnerability intersects with environmental and behavioral risks, accelerating disease onset and progression. This study emphasizes that effective management of adolescent diabetes requires more than clinical treatment alone. While advances in medical technologies and therapeutic approaches have improved glycemic control, long-term outcomes are strongly influenced by psychosocial support, family involvement, and access to comprehensive healthcare services. Integrated care models that combine medical, psychological, educational, and social interventions are essential for improving adherence, reducing complications, and enhancing quality of life. Furthermore, preventive strategies focusing on early screening, health education, and promotion of healthy lifestyles at the community and school levels are crucial to reducing disease burden. Strengthening medical and social care systems, particularly in resource-limited settings, will play a key role in addressing the rising prevalence of diabetes mellitus among adolescents and ensuring sustainable health outcomes.

## References

1. World Health Organization. Global report on diabetes. Geneva: WHO; 2016.
2. International Diabetes Federation. IDF Diabetes Atlas. 10th ed. Brussels: International Diabetes Federation; 2021.
3. Mayer-Davis EJ, Lawrence JM, Dabelea D, et al. Incidence trends of type 1 and type 2 diabetes among youths, 2002–2012. *New England Journal of Medicine*. 2017;376(15):1419–1429.
4. Rewers M, Ludvigsson J. Environmental risk factors for type 1 diabetes. *The Lancet*. 2016;387(10035):2340–2348.
5. Pinhas-Hamiel O, Zeitler P. The global spread of type 2 diabetes mellitus in children and adolescents. *Journal of Pediatrics*. 2005;146(5):693–700.
6. Pulgaron ER, Delamater AM. Obesity and type 2 diabetes in children: epidemiology and treatment. *Current Diabetes Reports*. 2014;14(8):508.
7. Hood KK, Rohan JM, Peterson CM, Drotar D. Interventions with adherence-promoting components in pediatric type 1 diabetes. *Diabetes Care*. 2010;33(7):1658–1664.
8. Silverstein J, Klingensmith G, Copeland K, et al. Care of children and adolescents with type 1 diabetes. *Diabetes Care*. 2005;28(1):186–212.
9. Cameron FJ, Garvey K, Hood KK. Adolescence and diabetes: challenges and opportunities. *The Lancet Diabetes & Endocrinology*. 2018;6(1):61–73.
10. UNICEF. Adolescent health and development: global challenges. New York: United Nations Children's Fund; 2020.

