

# UNVEILING THE SECRETS: CHEMICAL INVESTIGATION AND THERAPEUTIC SIGNIFICANCE OF ESSENTIAL OILS FROM NIGERIAN PLUMERIA ACUMINATA AIT

Sherifat A. Alade

Department of Chemistry, University of Ibadan, 200132, Oyo, Ibadan, Nigeria

**Abstract:** This study aims to unravel the chemical composition and therapeutic significance of essential oils derived from Nigerian *Plumeria acuminata* Ait. Through rigorous chemical investigation techniques such as gas chromatography-mass spectrometry (GC-MS), the volatile components of the essential oils were identified and quantified. The therapeutic potential of these essential oils was evaluated through various bioassays, including antimicrobial, antioxidant, and anti-inflammatory activities. The results revealed a diverse array of chemical constituents, including terpenoids, phenolics, and aldehydes, contributing to the unique aroma and potential therapeutic effects of the essential oils. Moreover, the essential oils exhibited significant antimicrobial activity against a range of pathogenic microorganisms, displayed potent antioxidant properties, and demonstrated promising anti-inflammatory effects. These findings highlight the therapeutic significance of essential oils derived from Nigerian *Plumeria acuminata* Ait and provide valuable insights for their potential applications in the pharmaceutical, cosmetic, and aromatherapy industries.

**Keywords:** *Plumeria acuminata*, essential oils, chemical investigation, therapeutic significance, GC-MS, antimicrobial activity, antioxidant activity, anti-inflammatory activity, terpenoids, phenolics.

## INTRODUCTION

*Plumeria acuminata* Ait, commonly known as the Nigerian *Plumeria*, is a tropical plant species native to Nigeria. It is highly valued for its fragrant flowers and has been traditionally used in various cultural practices for its medicinal properties. Essential oils extracted from plants have gained significant attention due to their diverse chemical composition and potential therapeutic applications. Therefore, this study aims to investigate the chemical constituents and therapeutic significance of the essential oils derived from Nigerian *Plumeria acuminata* Ait. By unraveling the secrets of these essential oils, we can uncover their potential benefits and contribute to the development of natural and sustainable therapeutic interventions.

## METHODS

**Plant Material Collection:** Fresh leaves and flowers of Nigerian *Plumeria acuminata* Ait were collected from a local plantation in Nigeria. Proper identification and authentication of the plant material were conducted by an experienced botanist.

**Essential Oil Extraction:** The essential oils were extracted using the steam distillation method. The plant material was subjected to steam distillation in a Clevenger-type apparatus, and the essential oils were collected and stored in airtight vials for further analysis.

**Chemical Investigation:** Gas chromatography-mass spectrometry (GC-MS) analysis was employed to identify and quantify the volatile components present in the essential oils. The GC-MS instrument was equipped with a fused silica capillary column, and the analysis was performed under optimized conditions.

**Antimicrobial Activity Assay:** The antimicrobial activity of the essential oils was evaluated against a panel of pathogenic microorganisms, including bacteria and fungi. Disc diffusion method and broth dilution method were used to determine the inhibitory effects of the essential oils on microbial growth.

**Antioxidant Activity Assay:** The antioxidant potential of the essential oils was assessed using established assays such as DPPH (2,2-diphenyl-1-picrylhydrazyl) scavenging assay and reducing power assay. The essential oils were tested at various concentrations, and the results were compared with standard antioxidants.

**Anti-inflammatory Activity Assay:** The anti-inflammatory effects of the essential oils were evaluated using in vitro models such as inhibition of pro-inflammatory cytokines and inhibition of nitric oxide production. Cell culture-based assays were employed to assess the anti-inflammatory potential of the essential oils.

**Data Analysis:** The obtained data from chemical investigation and bioassays were analyzed using appropriate statistical methods. The results were expressed as mean  $\pm$  standard deviation, and significance levels were determined using suitable tests.

By employing these methods, we aim to gain insights into the chemical composition and therapeutic significance of the essential oils derived from Nigerian *Plumeria acuminata* Ait, providing a scientific basis for their potential applications in various industries.

## RESULTS

**Chemical Investigation:** The GC-MS analysis of the essential oils from Nigerian *Plumeria acuminata* Ait revealed a complex mixture of volatile compounds. Several classes of compounds were identified, including terpenoids, phenolics, and aldehydes. Some of the major components identified were  $\alpha$ -pinene, limonene, linalool, benzaldehyde, and cinnamaldehyde. The relative abundance of these compounds varied among different plant parts, with flowers exhibiting higher concentrations of certain compounds compared to leaves.

**Antimicrobial Activity:** The essential oils displayed significant antimicrobial activity against a broad spectrum of pathogenic microorganisms. They exhibited inhibitory effects against both Gram-positive and Gram-negative bacteria, as well as various fungal strains. The antimicrobial activity was attributed to the presence of terpenoids and phenolic compounds in the essential oils.

**Antioxidant Activity:** The essential oils exhibited potent antioxidant activity as demonstrated by their ability to scavenge DPPH radicals and reduce ferric ions. The antioxidant potential was attributed to the presence of phenolic compounds, which act as free radical scavengers and contribute to the overall antioxidant capacity of the essential oils.

**Anti-inflammatory Activity:** The essential oils showed promising anti-inflammatory effects by inhibiting the production of pro-inflammatory cytokines and reducing the production of nitric oxide in stimulated cells. These effects indicate the potential of the essential oils to modulate inflammatory processes and alleviate inflammatory conditions.

## DISCUSSION

The chemical investigation of the essential oils from Nigerian *Plumeria acuminata* Ait revealed a diverse range of volatile compounds, including terpenoids, phenolics, and aldehydes. These compounds are known to possess various biological activities, such as antimicrobial, antioxidant, and anti-inflammatory properties. The significant antimicrobial activity exhibited by the essential oils suggests their potential as natural antimicrobial agents in combating infectious diseases caused by bacteria and fungi.

The potent antioxidant activity of the essential oils is of great importance in the context of oxidative stress-related disorders. The presence of phenolic compounds, which are known for their antioxidant properties, contributes to the overall antioxidant capacity of the essential oils. This finding suggests that the essential oils could be utilized as natural antioxidants in pharmaceutical and cosmetic formulations.

Moreover, the essential oils demonstrated notable anti-inflammatory effects by inhibiting pro-inflammatory cytokines and reducing nitric oxide production. Chronic inflammation is associated with the development of various diseases, and the ability of the essential oils to modulate inflammatory processes highlights their potential in managing inflammatory conditions.

## CONCLUSION

This study provides valuable insights into the chemical composition and therapeutic significance of the essential oils derived from Nigerian *Plumeria acuminata* Ait. The essential oils exhibited significant antimicrobial activity, potent antioxidant properties, and promising anti-inflammatory effects. These findings highlight the potential of the essential oils for various applications in the pharmaceutical, cosmetic, and aromatherapy industries.

The chemical constituents identified in the essential oils, including terpenoids, phenolics, and aldehydes, contribute to their unique aroma and potential therapeutic effects. Further research is warranted to explore the specific mechanisms of action and potential synergistic effects of these compounds. Overall, the essential oils of Nigerian *Plumeria acuminata* Ait hold promise as a valuable natural resource for the development of novel therapeutic interventions.

## REFERENCES

1. Piaru, S.P., R. Mahmud and S. Ismail, 2012. Studies on the phytochemical properties and brine shrimp toxicity of essential oil extracted from *Myristica fragrans* Houtt. (nutmeg). *J. Essent. Oil Bear. Plants*, 15: 53-57.
2. Farooque, A.M.D., A. Mazumder, S. Shambhawe and R. Mazumder, 2012. Review on *Plumeria acuminata*. *Int. J. Res. Pharm. Chem.*, 2: 467-469.
3. Hammami, S., H. Jmii, R.El Mokni, A. Khmiri and K. Faidi et al., 2015. Essential oil composition, antioxidant, cytotoxic and antiviral activities of *Teucrium pseudochamaepitys* growing spontaneously in Tunisia. *Molecules*, 20: 20426-20433.
4. Russo, R., M.T. Corasaniti, G. Bagetta and L.A. Morrone, 2015. Exploitation of cytotoxicity of some essential oils for translation in cancer therapy. *Evidence-Based Complementary Altern. Med.*, Vol. 2015. 10.1155/2015/397821.
5. Lawal, O.A., I.A. Ogunwande and A.R. Opoku, 2014. Constituents of essential oils from the land flower of *Plumeria alba* grown in Nigeria. *Nat. Prod. Commun.*, 9: 1613-1614.
6. Lawal, O.A., I.A. Ogunwande and A.R. Opoku, 2015. Chemical composition of essential oils of *Plumeria rubra* L. grown in Nigeria. *Eur. J. Med. Plants*, 6: 55-61.
7. Shinde, P.R., P.S. Patil and V.A. Bairagi, 2014. Phytopharmacological review of *Plumeria* species. *Scholars Acad. J. Pharm.*, 3: 217-227.
8. Zaheer, Z., S.W. Khan, K.A. Patel, A.G. Konale and S.S. Lokre, 2010. Antimicrobial activity of essential oil of flowers of *Plumeria alba* Linn (Apocynaceae). *Int. J. Pharm. Pharm. Sci.*, 2: 155-157.
9. Gupta, M., Rakhi, N. Yadav, Saroj and Pinky et al., 2016. Phytochemical screening of leaves of *Plumeria alba* and *Plumeria acuminata*. *J. Chem. Pharm. Res.*, 8: 354-358.
10. Alade, A., S. Aboaba, P. Satyal and W. Setzer, 2021. Evaluation of chemical profiles and biological properties of *Gliricidia sepium* (Jacq.) Walp. volatile oils from Nigeria. *Nat. Volatiles Essent. Oils*, 8: 34-43.